## **Asthma Action Plan for Home and School**



Name						DOB	/
1 <del>-</del>	n □ Intermittent □ Mild Pe					t	
Peak Flow Meter Per	sonal Best						
Green Zone: Doir	ng Well						
	ng is good - No cough or whe			Sleeps	well at night		
Control Medicine(s)	Medicine	How much t	SWA BRICHSWELL		and how often to tak		Take at ☐ Home ☐ School ☐ Home ☐ School
Physical Activity	☐ Use albuterol/levalbuterol						20 90 900 90 90 90 91 20
Yellow Zone: Cau	rtion						
6 6	roblems breathing – Cough, v ow Meterto(betv		1 <del>7</del> 4		orking or playing -	Wake at night	
Quick-relief Medicin Control Medicine(s)		e medicines					
$\Box$ Add $\_$ $\Box$ Change to $\_$ The child should feel better within 20–60 minutes of the quick-relief treatment. If the child is getting worse or is in the Yellow Zone for more than 24 hours, THEN follow the instructions in the RED ZONE and call the doctor right away!							
Red Zone: Get He	elp Now!						
Symptoms: Lots of problems breathing - Cannot work or play - Getting worse instead of better - Medicine is not helping  Peak Flow Meter (less than 50% of personal best)							
Take Quick-relief M	edicine NOW!	evalbuterol	puffs,			(how frequ	uently)
Call 911 immediatel	y if the following danger signs		<ul> <li>Lips or finger</li> </ul>	rnails are		of breath	
School Staff: Follow the Yellow and Red Zone instructions for the quick-relief medicines according to asthma symptoms.  The only control medicines to be administered in the school are those listed in the Green Zone with a check mark next to "Take at School".							
Healthcare Provider							
Name		Date	Phone ()		Signature		
☐ I consent to comm	or the medicines listed in the a unication between the prescri providers necessary for asthm	bing health car	e provider or o	clinic, the	e school nurse, the so		
Name		Date	Phone (	_	Signature		